



日本入国時に必要な検査証明書の要件（検体、検査方法、検査時間）

※ 有効な検体、検査方法等が記載された検査証明書のみ有効と取り扱います。

<有効な検査証明書として認められる要件>

検体

- Nasopharyngeal/Nasopharynx
(Swab/smear)
Rhinopharyngeal/Rhinopharynx
(Swab/Smear)
(鼻咽頭ぬぐい液)
- Nasal Swab (鼻腔ぬぐい液)
- (Deep throat) Saliva (唾液)
- Nasopharyngeal (※) (and /, /+)
oropharyngeal(throat) (swabs /smear)
(鼻咽頭ぬぐい液・咽頭ぬぐい液の混合)
(Naso and oropharyngeal/Rhino and oropharyngeal/oro and nasopharyngeal (※))
(※Nasopharyngeal/Nasopharynx/Rhinopharyngeal/Rhinopharynx)

<有効な検査証明書として認められない主な例>

- × Oral (swab/smear) (口腔ぬぐい)
- × Throat (swab/smear) (咽頭ぬぐい)
- × Mid-Turbinate Nasal (swab/smear) (中鼻甲介ぬぐい)
- × Gargle Water (うがい液)
- × mixture of sample "A" and "B"
(「鼻咽頭ぬぐい液と咽頭ぬぐい液の混合検体」を除く、複数箇所から採取した検体の混合検体)
(なお、“A”、“B”は検体を指す)

<Example>

- × Nasal and throat (swab/smear) (鼻腔・咽頭ぬぐい)
- × Pharyngeal and nasal (swab/smear) (咽頭・鼻腔ぬぐい)
- × Nasal and oropharyngeal/oropharynx (swab/smear)
(鼻腔・口腔咽頭ぬぐい)

検査方法

※鼻腔ぬぐい液検体は核酸増幅検査のみ有効

- ※下記の方法、もしくは検査手法名を含む方法
- Nucleic acid amplification test ((RT-) PCR)
核酸増幅検査 ((RT-) PCR法)
(real-time (RT-)PCR、Q-PCR、Fluorescence-PCR、Multiplex-PCR)
 - Nucleic acid amplification test (LAMP)
核酸増幅検査 (LAMP法) (RT-LAMP)
 - Nucleic acid amplification test (TMA)
核酸増幅検査 (TMA法)
 - Nucleic acid amplification test (TRC)
核酸増幅検査 (TRC法)
 - Nucleic acid amplification test (Smart Amp)
核酸増幅検査 (Smart Amp法)
 - Nucleic acid amplification test (NEAR)
核酸増幅検査 (NEAR法) (例: ID-NOW®)
 - Next generation sequence
次世代シーケンス法
 - Quantitative antigen test (CLEIA/ECLIA)
抗原定量検査

× Antigen (test/kit)
(抗原検査)

× Rapid antigen (test/kit)
(迅速抗原検査)

× Antibody (test/kit)
(抗体検査)

※日本で無症状者への検査として推奨している検体・検査方法。

※日本で無症状者への検査として推奨されていない検体・検査方法。

検査時間

- 検体採取が
出国前の72時間以内

- × 結果判明が
出国前の72時間以内

※今後、国内外の状況に鑑み、上記取扱いを変更する可能性があります。

参考：新型コロナウイルス感染症（COVID-19）病原体検査の指針第5版 (<https://www.mhlw.go.jp/content/000788513.pdf>)



Requirements for Certificate of Testing for Entering Japan

※Only test certificates which show the results of the test using valid samples and testing methods below are valid.

Valid

Invalid

Sample:

- Nasopharyngeal/Nasopharynx (Swab/smear)
- Rhinopharyngeal/Rhinopharynx (Swab/smear)
- Nasal Swab
- (Deep throat)Saliva sample
- Nasopharyngeal (※) (and /, /+) oropharyngeal(throat) (swab /smear)
(Naso and oropharyngeal/Rhino and oropharyngeal/oro and nasopharyngeal (※))
(※Nasopharyngeal/Nasopharynx/Rhinopharyngeal/Rhinopharynx)

※ Nasal Swab is valid when the test method is Nucleic acid amplification test.

- Oral (swab/smear)
- Throat (swab/smear)
- Mid-Turbinate Nasal (swab/smear)
- Gargle Water
- Mixture of sample “A” and “B”
“A” and “B” show all types of sample(except nasopharyngeal and oropharyngeal(swab/smear)).
<Example>
 - Nasal and throat (swab/smear)
 - Pharyngeal and nasal (swab/smear)
 - Nasal and oropharyngeal/oropharynx (swab/smear)

Testing Method:

Nucleic acid amplification test

- PCR - Polymerase chain reaction
- (RT-)PCR - Reverse transcription PCR (real-time (RT-)PCR/Q-PCR/Fluorescence-PCR/Multiplex-PCR)
- LAMP – Loop-mediated isothermal amplification (RT-LAMP)
- TMA – Transcription mediated amplification
- TRC - Transcription-reverse transcription concerted reaction
- Smart Amp - Smart amplification process
- NEAR - Nicking endonuclease amplification reaction (e.g. ID-NOW)

- Antigen test/test kit
- Rapid antigen test/test kit
- Antibody test/test kit

Others

- Next generation sequencing
- Quantitative antigen test (CLEIA/ECLIA)

※Recommend in Japan as a sample/testing method for asymptomatic people.

※Not recommend in Japan as a sample/testing method for asymptomatic people.

Sample collection must be done Within 72 hours before the departure time of the flight.

※These requirements may be changed depending on infection status.